

THE LANCET

Infectious Diseases

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed.
We post it as supplied by the authors.

Supplement to: Rajasingham R, Smith RM, Park BJ, et al. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet Infect Dis* 2017; published online May 5. [http://dx.doi.org/10.1016/S1473-3099\(17\)30243-8](http://dx.doi.org/10.1016/S1473-3099(17)30243-8).


Appendix

Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning DW, Loyse A, Boulware DR. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet Infectious Diseases*. 2017.

Table of Contents


Supplemental Figure 1: Methodology of calculations	2
Supplemental Methods.....	4
Supplemental Table 1: Estimation and Uncertainty Model Inputs	6
Supplemental Figure 2: Cryptococcal Meningitis Annual Incidence	7
Supplemental Table 2: Cryptococcal Antigen Prevalence Rates.....	8
References	12

Supplemental Figure 1: Methodology of calculations of those at risk for cryptococcal infection, CRAG+, number who develop cryptococcal meningitis, and number who die of cryptococcal meningitis.



Notes: LTFU = Lost to follow up; VF = viral failure; LMICs = low and middle-income countries; HICs = high-income countries.
 CD4 100-200 calculated as above though percentage of CD4<350 is 27.5% in LMICs, and 31.5% in HICs (refer to Table 1)
 Boxed letters correspond to Table 1

At risk for Cryptococcal infection X CRAG Prevalence



Supplemental Methods

Global Burden of HIV Infection

Details regarding the assumptions behind the UNAIDS uncertainty bounds are published.¹ Where possible we used UNAIDS and WHO reports and multinational studies, as these estimates were less biased than single-center country-specific research publications. However, for many high-income countries in Europe and North America no estimates were made by UNAIDS or WHO of adults eligible for ART, receiving ART, or percent coverage. As available, UNAIDS Country Progress reports were used for Canada and China. U.S. HIV statistics were taken from CDC reports.² Regional estimates for Africa, Asia and Pacific, Caribbean, Latin America, and Western Pacific were simply the sum of each country's UNAIDS estimate. Notably, Japan, Korea, and New Zealand were not included in the Asian estimate, due to lack of accurate data. UNAIDS regional estimates were used for Europe, Eastern Mediterranean and North African region given that many country-specific estimates were unavailable.

CRAG Prevalence

For countries with available data, CRAG prevalence amongst persons with a CD4<100 cells/ μ L was used for that specific country along with the associated 95%CI. If the source cited CRAG prevalence for those with CD4>100 cells/ μ L, raw numbers were analyzed as possible to calculate CRAG prevalence and confidence interval for CD4<100 cells/ μ L. If raw numbers were not available, we still included the study in this analysis. If more than one study existed for a country, a weighted average with 95%CI was calculated, based on study size. For countries without specific published data, a weighted mean of the CRAG prevalence for the region with 95%CI was used as an estimate. As there were no studies published in the Caribbean, a weighted average of CRAG prevalence from Latin America was used. All literature cited refers to *Cryptococcus neoformans*, not *gattii* infection.

There is a small risk of cryptococcal disease amongst those with CD4 >100 cells/ μ L.³⁻⁵ We estimated the number with a CD4 between 100-200 cells/ μ L as 27.5% (24% to 31%) of those with CD4<350 cells/ μ L in LMICs. In HICs we estimated 31.5% (29% to 34%) of those with a CD4<350cells/ μ L had a CD4 between 100 and 200 cells/ μ L.^{6,7} In persons with CD4 100-200 cells/ μ L, we estimated a CRAG prevalence of one fourth of the prevalence amongst those with a CD4<100 cells/ μ L, based on a weighted average of CRAG prevalence in those with CD4>100 cells/ μ L compared with CD4<100 cells/ μ L.^{3-5,8-18}

CRAG-positive persons who develop Cryptococcal Meningitis

Estimates of the number of CRAG-positive adults who go on to develop symptomatic cryptococcal meningitis were from published cohort studies,^{3,4,10,19,20} where 33% (95%CI: 25% to 41%) of CRAG-

positive persons had prevalent disease at time of screening or developed *known* cryptococcal meningitis after initiating ART. In those research cohorts, 60% (72/121) died (95%CI: 50% to 68%).^{3,4,10,19,20} Notably, these research studies reported a short time from CRAG positivity to ART initiation (≤ 2 weeks), and all of them were performed in sub-Saharan Africa or Asia. In routine care, there is likely further delay in initiating ART, and thus increased disease progression to meningitis. However, ART-mediated immune recovery may unmask the initially subclinical disseminated cryptococcal infection resulting in non-meningitis disease presentations (e.g. sepsis, pneumonitis).^{3,21} From these studies, we presumed that a significant proportion of those CRAG+ who died of unknown causes or were lost to follow up in these studies were in fact deaths related to cryptococcosis. Including those who died or were lost to follow up as presumed cryptococcosis, we estimated 70% (Range: 56% to 82%) of CRAG+ persons developed cryptococcal disease or died despite initiating ART, unless preemptively treated. Without ART or preemptive fluconazole therapy, we assumed 95% (Range: 90% to 100%) progression in all regions (consensus expert opinion of authors).^{22,23}

Cryptococcal Meningitis Deaths

We used UNAIDS country-level data on ART access as the surrogate for the proportion with access to quality medical care. We estimated 1-year mortality in low-income countries (LICs) as 70% (95%CI: 59% to 81%) after cryptococcal meningitis for those in care and 100% for those not in care.^{30, 47-52} Uncertainty ranges were calculated by pooling the data from each paper, and calculating the 95% confidence interval. Data from research clinical trials suggest 10-week mortality of 39% (95%CI: 36% to 42%), with additional ~15% mortality through 1-year.^{9,24,25} Outside of research trials, we expected mortality to be higher.²² Many in LICs lack an initial diagnosis, receive fluconazole monotherapy, and do not receive therapeutic lumbar punctures, thus 70% mortality may be optimistic.^{9,25-32} All sub Saharan African countries were considered low-income countries for the purpose of this estimate. Despite South Africa and Botswana having a higher GDP than most other African nations, the rate of CRAG+ progression and mortality from cryptococcal meningitis in these countries is similar to those of low-income nations.

Uncertainty Analysis


When data were lacking, we used expert opinion to define distributions, based on previously published estimates. Specifically, expert opinion of the authors was used to estimate progression from CRAG+ to cryptococcal meningitis for those not in care, as for those not in care there are no published estimates in the literature. Deaths from cryptococcal meningitis for those not in care were also estimated this way.

Supplemental Table 1: Estimation and Uncertainty Model Inputs

Model Input	Data Source(s)	Distribution	Parameters
Adults living with HIV	UNAIDS 2013 estimate: 31.8 million		Published low, high values: 30.1 million, 33.7 million
Adults with CD4<350 cells/ μ L	UNAIDS 2013 estimate of those eligible for ART (i.e. CD4<350 cells/ μ L): 19.5 million		Low, high values: 18.5 million, 20.7 million – from UNAIDS 2016 range
Adults receiving ART	UNAIDS 2013 estimate 11.3 million		Published low, high values: 10.4 million, 12.7 million
New initiation of ART	UNAIDS 2013, 2014 of Receipt of ART minus LTFU		Low, high values: 1.8 million, 2.8 million (80% and 120% of estimate)
CD4<100 cells/ μ L (LMICs)	22.5% of those with CD4<350 cells/ μ L ^{6,7,33-35}	Beta	Low, high values: 19%, 26%;
CD4<100 cells/ μ L (HICs)	18.5% of those with CD4<350 cells/ μ L ^{6,7,33-35}	Beta	Low, high values: 16% to 21%;
CD4 100 to 200 cells/ μ L (LMICs)	27.5% of those with CD4<350 cells/ μ L ^{6,7,33-35}	Beta	Low, high values: 24% to 31%;
CD4 100 to 200 cells/ μ L (HICs)	31.5% of those with CD4<350 cells/ μ L ^{6,7,33-35}	Beta	Low, high values: 29% to 34%;
Virologic failure (within first year of ART)	16% of those on ART	Beta	Low, high values: 12% to 20%;
Virologic failure (after 1 year on ART)	5.33% of those on ART	Beta	Low, high values: 4.0% to 6.6%;
CD4 100 to 200 cells/ μ L not on ART, LTFU, or virologic failure	25% of those with CD4<100 not on ART, or LTFU, or with virologic failure ^{3,12}	Beta	N/A
CRAG prevalence for countries without primary data	Regional weighted average	Normal	Computed 95% CI
CRAG+ who develop meningitis			
-- In care without preemptive treatment	70% ^{3,4,10,19,20}	Beta	Low value, high value: 56%, 84%;
-- Not in care	95% ^{3,4,10,19,20}	Beta	Low value, high value: 90%, 100%;
CRAG negative progression to CRAG+ without ART (LMICs)	5.1% annual incidence with 50% competing risk for starting ART or death	Beta	Low value, high value: 2.6%, 9.0%;
CRAG negative progression to CRAG+ without ART (HICs)	2.55% annual incidence with 50% competing risk for starting ART or death	Beta	Low value, high value: 0.1%, 1.0%;
1-year mortality from meningitis (LICs)	70% ^{9,25,28-32}	Beta	Low value, high value: 59%, 81%;
1-year mortality from meningitis (MICs)	40% ^{9,36-38}	Beta	Low value, high value: 34%, 46%;
1-year mortality from meningitis (North America)	20% ^{39,40}	Beta	Low value, high value: 12.5%, 27.5%;
1-year mortality from meningitis (Europe)	30% (including Ukraine, Russia)	Beta	Low value, high value: 25%, 35%;
Not in care (all regions)	1.5x higher ²²		N/A

LTFU =Lost to follow up; LIC = low income country; MIC = middle income country. References numbers as in the manuscript.

Supplemental Figure 2: Cryptococcal meningitis annual incidence. Annually, an estimated 223,100 (95% CI, 150,600 to 282,400) cryptococcal meningitis cases occur globally.



Supplemental Table 2: Cryptococcal Antigen Prevalence Rates (Source Data for Figure 1)

<i>Year</i>	<i>Author</i>	<i>Country</i>	<i>Setting</i>	<i>Overall prevalence</i>	<i>Numerator</i>	<i>Denominator</i>	<i>95%CI Lower</i>	<i>95%CI Higher</i>	<i>HIV severity</i>
1989	Desmet ⁴¹	Dem. Rep. of the Congo	Newly diagnosed HIV+ patients	12.2%	55	450	9.3%	15.6%	Unknown
1992	Swinne ⁴²	Rwanda	Random HIV+ sera	4.2%	9	213	2.0%	7.9%	Unknown
1995	Negroni ⁴³	Argentina	Unknown	6.7%	13	193	3.6%	11.2%	193 CD4 <300
2003	Tassie ⁴⁴	Uganda (Mbarara)	Inpatient (n=144) and outpatient (n=53) HIV+	4.4%	8	182	1.9%	8.5%	Stage III or IV
2007	Liechty ⁴⁵	Uganda (Tororo)	HIV+ initiating ART	5.8%	22	377	3.7%	8.7%	CD4 <100
2007	Micol ⁴⁶	Cambodia	HIV+ seen at hospital associated ART programs	10.8%	32	295	7.5%	15.0%	CD4 <200
2009	Jarvis ¹⁹	South Africa (Cape Town)	HIV+ initiating ART	6.7%	21	312	4.2%	10.1%	CD4 <100
2010	Meya ³	Uganda (Kampala)	HIV+ initiating ART	8.8%	26	295	5.8%	12.6%	CD4 <100
2010	Pongsai ¹⁰	Thailand (Bangkok)	HIV+, ART-naïve	12.9%	11	85	6.6%	22.0%	CD4 <100
2011	Parkes-Ratanshi ¹²	Uganda (Masaka)	HIV+, ART-naïve adults	7.8%	59	757	6.0%	9.9%	CD4 <100
2011	Mamoojee ⁴⁷	Ghana	HIV+ initiating ART	2.0%	2	92	0.3%	7.6%	CD4 <100
2011	Kendi ⁴⁸	Kenya	HIV+ initiating ART	7.2%	66	920	5.6%	9.0%	CD4 ≤100
2012	Osazuwa ¹¹	Nigeria	HIV+ initiating ART	12.7%	19	150	7.8%	19.1%	CD4 <200
2012	Linares ⁴⁹	Peru (Lima)	HIV+, ART-naïve	3.6%	13	365	1.9%	6.0%	CD4 ≤100
2012	Kwan ¹³	Thailand	HIV+ women initiating ART	11.0%	9	84	5.0%	19.4%	CD4 <100
2012	Bedell ⁵⁰	Malawi	HIV+ initiating ART	1.7%	8	469	0.7%	3.3%	All (median=129)

2013	Alemu ¹⁴	Ethiopia (Addis)	HIV+, both ART-naïve and ART-experienced	11.2%	13	116	6.1%	18.4%	CD4 <100
2013	Patel ⁵¹	U.K. (London)	Newly diagnosed HIV+ patients	5.0%	8	157	2.2%	9.8%	CD4 <100
2013	Meyer ⁴⁸	Kenya (multisite)	HIV+ initiating ART	9.1%	117	1286	7.6%	10.8%	CD4 ≤100
2013	Escandón ⁵²	Colombia	ART-naïve HIV+	7.1%	21	297	4.4%	10.6%	All
2013	Smith ⁵³	Vietnam (multisite)	HIV+ initiating ART	4.0%	9	226	1.8%	7.4%	CD4 <100
2013	Chukwuanukwu ⁵⁴	Nigeria	Pregnant HIV+	13.0%	21	160	8.3%	19.4%	All
2013	Rugemalila ¹⁵	Tanzania (Moshi)	Outpatient HIV+ initiating ART or on ART<6 months	4.8%	6	124	1.8%	10.2%	CD4 <100
2013	Beyene ⁵	Ethiopia (Oromia Region)	HIV+ initiating ART-naïve and defaulters	15.5%	16	103	9.1%	24.0%	All
2013	Costa ¹⁷	Brazil (Para State)	HIV+ in 2 HIV clinics on ART	2.6%	11	418	1.3%	4.7%	All
2014	McKenney ⁵⁵	U.S. (multisite)	Stored sera of HIV+ both ART-naïve and ART-experienced	2.9%	54	1,872	2.2%	3.7%	CD4 <100
2014	Drain ⁵⁶	South Africa (Durban)	HIV+ initiating ART in 4 outpatient sites, urine LFA	8.7%	67	773	6.8%	10.9%	CD4 ≤100
2014	Ganiem ⁵⁷	Indonesia	Stored sera of HIV+ ART-naïve outpatients in 1 clinic using LFA	7.1%	58	810	5.5%	9.2%	CD4 <100
2014	Sawadogo ⁷	Namibia	Prospective CD4	4.1%	21	516	2.5%	6.2%	CD4 <100
2014	MSF ⁵⁸	Dem. Rep. of the Congo	HIV Outpatient Clinic	15.2%	93	613	12.4%	18.3%	CD4 <100
2014	MSF ⁵⁸	Guinea	HIV Outpatient Clinic	4.9%	4	82	1.3%	12.0%	CD4 <100
2014	MSF ⁵⁸	Mozambique	HIV Tertiary Referral Clinic	6.6%	11	166	3.4%	11.5%	CD4 <100

2014	MSF ⁵⁸	Lesotho	HIV Outpatient Clinic	5.4%	3	56	1.1%	14.9%	CD4 <100
2014	MSF ⁵⁸	Kenya (Kibera)	HIV Outpatient Clinic	12.8%	17	133	7.6%	19.7%	CD4 <100
2014	MSF ⁵⁸	Zimbabwe (Gutu, Buhera)	HIV Outpatient Clinic	7.1%	32	448	4.9%	9.9%	CD4 <100
2014	Magambo ⁵⁹	Tanzania (Mwanza)	HIV Outpatient Clinic	8.2%	6	73	3.1%	17.0%	CD4 <100
2015	Letang ⁴	Tanzania (Ifakara)	Stored plasma samples of HIV+ ART-naïve patients	4.3%	24	556	2.8%	6.4%	CD4 <100
2015	Govender ⁶⁰	South Africa (Gauteng)	HIV+ patients of 26 health facilities	4.0%	60	1494	3.1%	5.1%	CD4 <100
2015	Chipungu ⁵⁵	Malawi	Outpatient CD4<100	3.5%	2	57	0.4%	12.1%	CD4 <100
2015	Mfinanga ⁶¹	Tanzania, Zambia	Outpatient CD4<100	4.6%	33	717	3.2%	6.4%	CD4 <100
2015	Pac ⁶²	Uganda	Outpatient CD4<100	6.8%	12	177	3.6%	11.5%	CD4<100
2016	Morawski ⁶³	Uganda (Multisite)	Outpatient CD4<100	8.7%	162	1860	7.5%	10.1%	CD4 <100
2016	Longley ⁶⁴	South Africa	Outpatient CD4<100	4.3%	28	645	2.9%	6.2%	CD4 <100
2016	Oladele ⁶⁵	Nigeria	Outpatient CD4<250	9.8%	6	61	4%	20%	CD4 <100
2016	Ezeanolue ⁶⁶	Nigeria	Outpatients CD4<100	3.7%	52	1399	2.8%	4.8%	CD4<100
2016	Vallabhaneni ⁶⁷	South Africa	Outpatients CD4<100	2.1%	24	1170	1.3%	3.0%	CD4<100
2016	Ogouyemiz ⁶⁸	Benin	Outpatients CD4<100	3.9%	6	154	1.0%	8.0%	CD4<200
2017	Kadam D ⁶⁹	India	Outpatient CD4<100	6.1%	11	180	3.1%	10.7%	CD4 <100
Total			Outpatients Only	6.4%	1365	21487	6.0%	6.7%	
Total			Outpatients LMIC	6.7%	1303	19458	6.3%	7.1%	

Studies of Primarily Hospitalized Populations not included in Figure 1.

<i>Year</i>	<i>Author</i>	<i>Country</i>	<i>Setting</i>	<i>Overall prevalence</i>	<i>Numerator</i>	<i>Denominator</i>	<i>95%CI Lower</i>	<i>95%CI Higher</i>	<i>HIV severity</i>
1997	Grant ⁷⁰	Ivory Coast	Inpatient HIV+	2.5%	5	199	0.8%	5.8%	All
2008	Oumar ⁷¹	Mali	Inpatient HIV+ patients in 1 ID unit hospital	3.1%	4	115	1.0%	8.7%	All
2011	Wajanga ⁷²	Tanzania	HIV+ inpatients	5.1%	17	333	3.0%	8.0%	All
2011	Apetse ⁷³	Togo	HIV+ inpatient 22 health sites India Ink, CrAg latex agglutination	2.9%	21	714	1.8%	4.5%	All
2012	Oyella ⁷⁴	Uganda	HIV+ inpatients (97%) and outpatients (3%)	19%	70	367	15.2%	23.5%	CD4 < 100
2012	Dzoyem ⁷⁵	Cameroon	Inpatient HIV+						
2013	Luma ⁷⁶	Cameroon	HIV+ patients	11.2%	75	672	8.9%	13.8%	All
2014	Okome-Nkoumou ⁷⁷	Gabon	HIV+ ART-naïve patients in 1 ID unit	0.44%	2	458	0.1%	1.6%	All
2015	Katchanov ⁷⁸	Germany	Hospital Admissions CD4<100	1.6%	28	1723	1.1%	2.3%	CD4 <100
2015	Frola ⁷⁹	Peru (Lima)	CD4<100 hospital	8.8%	10	114	4.3%	15.5%	CD4 <100
2016	Vidal ⁸⁰	Brazil	Inpatients CD4<200	3.1%	5	163	1.0%	7.0%	CD4 <200

References:

1. UNAIDS. Gap Report. 2014. www.unaids.org/en/resources/campaigns/2014gapreport (accessed 21 Dec 2015).
2. Bradley H, Hall HI, Wolitski RJ, et al. Vital Signs: HIV diagnosis, care, and treatment among persons living with HIV--United States, 2011. *MMWR Morb Mortal Wkly Rep* 2014; **63**(47): 1113-7.
3. Meya DB, Manabe YC, Castelnuevo B, et al. Cost-effectiveness of serum cryptococcal antigen screening to prevent deaths among HIV-infected persons with a CD4+ cell count < or = 100 cells/microL who start HIV therapy in resource-limited settings. *Clin Infect Dis* 2010; **51**(4): 448-55.
4. Letang E, Muller MC, Ntamatungiro AJ, et al. Cryptococcal antigenemia in immunocompromised HIV patients in rural Tanzania: A preventable cause of early mortality. *Open Forum Infect Dis* 2015; **2**(2): ofv046.
5. Beyene T, Woldeamanuel Y, Asrat D, Ayana G, Boulware DR. Comparison of cryptococcal antigenemia between antiretroviral naive and antiretroviral experienced HIV positive patients at two hospitals in Ethiopia. *PLoS One* 2013; **8**(10): e75585.
6. UNAIDS. Access to Antiretroviral Therapy in Africa: Status report on progress towards the 2015 targets. Geneva 2013: www.unaids.org/en/resources/documents/2013.
7. Avila D, Althoff KN, Mugglin C, et al. Immunodeficiency at the start of combination antiretroviral therapy in low-, middle-, and high-income countries. *J Acquir Immune Defic Syndr* 2014; **65**(1): e8-16.
8. Boulware DR, Meya DB, Muzoora C, et al. Timing of antiretroviral therapy after diagnosis of cryptococcal meningitis. *N Engl J Med* 2014; **370**(26): 2487-98.
9. Jarvis JN, Bicanic T, Loyse A, et al. Determinants of mortality in a combined cohort of 501 patients with HIV-associated Cryptococcal meningitis: implications for improving outcomes. *Clin Infect Dis* 2014; **58**(5): 736-45.
10. Pongsai P, Atamasirikul K, Sungkanuparph S. The role of serum cryptococcal antigen screening for the early diagnosis of cryptococcosis in HIV-infected patients with different ranges of CD4 cell counts. *J Infect* 2010; **60**(6): 474-7.
11. Osazuwa F, Dirisu JO, Okuonghae PE, Ugbebor O. Screening for cryptococcal antigenemia in anti-retroviral naive AIDS patients in benin city, Nigeria. *Oman Med J* 2012; **27**(3): 228-31.
12. Parkes-Ratanshi R, Wakeham K, Levin J, et al. Primary prophylaxis of cryptococcal disease with fluconazole in HIV-positive Ugandan adults: a double-blind, randomised, placebo-controlled trial. *Lancet Infect Dis* 2011; **11**(12): 933-41.
13. Kwan CK, Leelawiwat W, Intalapaporn P, et al. Utility of cryptococcal antigen screening and evolution of asymptomatic cryptococcal antigenemia among HIV-infected women starting antiretroviral therapy in Thailand. *J Int Assoc Provid AIDS Care* 2014; **13**(5): 434-7.
14. Alemu AS, Kempker RR, Tenna A, et al. High prevalence of cryptococcal antigenemia among HIV-infected patients receiving antiretroviral therapy in Ethiopia. *PLoS One* 2013; **8**(3): e58377.

15. Rugemalila J, Maro VP, Kapanda G, Ndaro AJ, Jarvis JN. Cryptococcal antigen prevalence in HIV-infected Tanzanians: a cross-sectional study and evaluation of a point-of-care lateral flow assay. *Trop Med Int Health* 2013; **18**(9): 1075-9.
16. Smitson CC, Tenna A, Tsegaye M, et al. No association of cryptococcal antigenemia with poor outcomes among antiretroviral therapy-experienced HIV-infected patients in Addis Ababa, Ethiopia. *PLoS One* 2014; **9**(1): e85698.
17. Costa MM, Madeira L, Feitosa RN, et al. Detection of *Cryptococcus neoformans* capsular antigen in HIV-infected patients in the state of Para in the north of Brazil. *Curr HIV Res* 2013; **11**(8): 647-51.
18. Magambo KA, Kalluvya SE, Kapoor SW, et al. Utility of urine and serum lateral flow assays to determine the prevalence and predictors of cryptococcal antigenemia in HIV-positive outpatients beginning antiretroviral therapy in Mwanza, Tanzania. *J Int AIDS Soc* 2014; **17**: 19040.
19. Jarvis JN, Lawn SD, Vogt M, Bangani N, Wood R, Harrison TS. Screening for cryptococcal antigenemia in patients accessing an antiretroviral treatment program in South Africa. *Clin Infect Dis* 2009; **48**(7): 856-62.
20. Govender N, Roy M, Mendes J, Zulu T, Chiller T, Karstaedt A. Evaluation of screening and treatment of cryptococcal antigenaemia among HIV-infected persons in Soweto, South Africa. *HIV Med* 2015.
21. Jacob ST, Moore CC, Banura P, et al. Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS ONE* 2009; **4**(11): e7782.
22. Mwaba P, Mwansa J, Chintu C, et al. Clinical presentation, natural history, and cumulative death rates of 230 adults with primary cryptococcal meningitis in Zambian AIDS patients treated under local conditions. *Postgrad Med J* 2001; **77**(914): 769-73.
23. Mayanja-Kizza H, Oishi K, Mitarai S, et al. Combination therapy with fluconazole and flucytosine for cryptococcal meningitis in Ugandan patients with AIDS. *Clin Infect Dis* 1998; **26**(6): 1362-6.
24. Boulware DR, Meya DB, Bergemann TL, et al. Clinical features and serum biomarkers in HIV immune reconstitution inflammatory syndrome after cryptococcal meningitis: a prospective cohort study. *PLoS Med* 2010; **7**(12): e1000384.
25. Butler EK, Boulware DR, Bohjanen PR, Meya DB. Long term 5-year survival of persons with cryptococcal meningitis or asymptomatic subclinical antigenemia in Uganda. *PLoS One* 2012; **7**(12): e51291.
26. Rolfes MA, Hullsiek KH, Rhein J, et al. The effect of therapeutic lumbar punctures on acute mortality from cryptococcal meningitis. *Clin Infect Dis* 2014; **59**(11): 1607-14.
27. Kambugu A, Meya DB, Rhein J, et al. Outcomes of cryptococcal meningitis in Uganda before and after the availability of highly active antiretroviral therapy. *Clin Infect Dis* 2008; **46**(11): 1694-701.
28. Gaskell KM, Rothe C, Gnanadurai R, et al. A prospective study of mortality from cryptococcal meningitis following treatment induction with 1200 mg oral fluconazole in Blantyre, Malawi. *PLoS One* 2014; **9**(11): e110285.
29. Lessells RJ, Mutevedzi PC, Heller T, Newell ML. Poor long-term outcomes for cryptococcal meningitis in rural South Africa. *S Afr Med J* 2011; **101**(4): 251-2.

30. Baldassarre R, Mdodo R, Omonge E, et al. Mortality after clinical management of AIDS-associated cryptococcal meningitis in Kenya. *East Afr Med J* 2014; **91**(5): 145-51.
31. Rothe C, Sloan DJ, Goodson P, et al. A prospective longitudinal study of the clinical outcomes from cryptococcal meningitis following treatment induction with 800 mg oral fluconazole in Blantyre, Malawi. *PLoS One* 2013; **8**(6): e67311.
32. Kendi C, Penner J, Koech J, et al. Predictors of outcome in routine care for Cryptococcal meningitis in Western Kenya: lessons for HIV outpatient care in resource-limited settings. *Postgrad Med J* 2013; **89**(1048): 73-7.
33. late presenters working group in COHERE in EuroCoord. Late presentation for HIV care across Europe: update from the Collaboration of Observational HIV Epidemiological Research Europe (COHERE) study, 2010 to 2013. *Euro Surveill* 2015; **20**(47).
34. European Centre for Disease Prevention and Control, WHO Regional Office for Europe. HIV/AIDS Surveillance in Europe, 2014. 2015.
<http://ecdc.europa.eu/en/healthtopics/aids/surveillance-reports/Pages/surveillance-reports.aspx> (accessed 1 Dec 2015).
35. Centers for Disease Control and Prevention. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 dependent areas—2013. *HIV Surveillance Supplemental Report 2015* 2015; **20**(2).
36. Rajasingham R, Rolfes MA, Birkenkamp KE, Meya DB, Boulware DR. Cryptococcal meningitis treatment strategies in resource-limited settings: a cost-effectiveness analysis. *PLoS Med* 2012; **9**(9): e1001316.
37. Vidal JE, Penalva de Oliveira AC, Dauar RF, Boulware DR. Strategies to reduce mortality and morbidity due to AIDS-related cryptococcal meningitis in Latin America. *Braz J Infect Dis* 2013; **17**(3): 353-62.
38. Jongwutiwes U, Kiertiburanakul S, Sungkanuparph S. Impact of antiretroviral therapy on the relapse of cryptococcosis and survival of HIV-infected patients with cryptococcal infection. *Curr HIV Res* 2007; **5**(3): 355-60.
39. Pyrgos V, Seitz AE, Steiner CA, Prevots DR, Williamson PR. Epidemiology of cryptococcal meningitis in the US: 1997-2009. *PLoS One* 2013; **8**(2): e56269.
40. Lortholary O, Poizat G, Zeller V, et al. Long-term outcome of AIDS-associated cryptococcosis in the era of combination antiretroviral therapy. *AIDS* 2006; **20**(17): 2183-91.
41. Desmet P, Kayembe KD, De Vroey C. The value of cryptococcal serum antigen screening among HIV-positive/AIDS patients in Kinshasa, Zaire. *AIDS* 1989; **3**(2): 77-8.
42. Swinne D, Bogaerts J, Van de Perre P, Batungwanayo J, Taelman H. Evaluation of the cryptococcal antigen test as a diagnostic tool of AIDS-associated cryptococcosis in Rwanda. *Ann Soc Belg Med Trop* 1992; **72**(4): 283-8.
43. Negroni R, Cendoya C, Arechavala AI, et al. Detection of *Cryptococcus neoformans* capsular polysaccharide antigen in asymptomatic HIV-infected patients. *Rev Inst Med Trop Sao Paulo* 1995; **37**(5): 385-9.
44. Tassie JM, Pepper L, Fogg C, et al. Systematic screening of cryptococcal antigenemia in HIV-positive adults in Uganda. *J Acquir Immune Defic Syndr* 2003; **33**(3): 411-2.

45. Liechty CA, Solberg P, Were W, et al. Asymptomatic serum cryptococcal antigenemia and early mortality during antiretroviral therapy in rural Uganda. *Trop Med Int Health* 2007; **12**(8): 929-35.
46. Micol R, Lortholary O, Sar B, et al. Prevalence, determinants of positivity, and clinical utility of cryptococcal antigenemia in Cambodian HIV-infected patients. *J Acquir Immune Defic Syndr* 2007; **45**(5): 555-9.
47. Mamoojee Y, Shakoor S, Gorton RL, et al. Short Communication: Low seroprevalence of cryptococcal antigenaemia in patients with advanced HIV infection enrolling in an antiretroviral programme in Ghana. *Trop Med Int Health* 2011; **16**(1): 53-6.
48. Meyer AC, Kendi CK, Penner JA, et al. The impact of routine cryptococcal antigen screening on survival among HIV-infected individuals with advanced immunosuppression in Kenya. *Trop Med Int Health* 2013; **18**(4): 495-503.
49. Linares Barandiaran LP, J.; Bustamante, B. . Cryptococcal antigenemia in Peruvian HIV-infected patients with a CD4 cell count <100 cell/mm³. 18th Congress of the International Society for Human and Animal Mycology 2012 (ISHAM2012); 2012 11–15 June 2012; Berlin, Germany; 2012.
50. Bedell RA, Anderson ST, van Lettow M, et al. High prevalence of tuberculosis and serious bloodstream infections in ambulatory individuals presenting for antiretroviral therapy in Malawi. *PLoS One* 2012; **7**(6): e39347.
51. Patel S, Shin GY, Wijewardana I, et al. The prevalence of cryptococcal antigenemia in newly diagnosed HIV patients in a Southwest London cohort. *J Infect* 2013; **66**(1): 75-9.
52. Escandon P, Lizarazo J, Agudelo CI, Chiller T, Castaneda E. Evaluation of a rapid lateral flow immunoassay for the detection of cryptococcal antigen for the early diagnosis of cryptococcosis in HIV patients in Colombia. *Med Mycol* 2013; **51**(7): 765-8.
53. Smith RM, Nguyen TA, Ha HT, et al. Prevalence of cryptococcal antigenemia and cost-effectiveness of a cryptococcal antigen screening program--Vietnam. *PLoS One* 2013; **8**(4): e62213.
54. Chukwuanukwu R MP, Iloghalu E, Onyenekwe C, Ifeanyichukwu M, Mbamalu C. *Cryptococcus neoformans* Antigenemia in HIV Positive Pregnant Women Attending a PMTCT Clinic in South-East Nigeria. *Journal of Biology, Agriculture and Healthcare* 2013; **3**(18): 15-20.
55. McKenney J, Bauman S, Neary B, et al. Prevalence, correlates, and outcomes of cryptococcal antigen positivity among patients with AIDS, United States, 1986-2012. *Clin Infect Dis* 2015; **60**(6): 959-65.
56. Drain PK, Kleene JM, Coleman SM, et al. Prevalence of cryptococcal antigenuria at initial HIV diagnosis in KwaZulu-Natal. *HIV Med* 2015; **16**(10): 640-4.
57. Ganiem AR, Indrati AR, Wisaksana R, et al. Asymptomatic cryptococcal antigenemia is associated with mortality among HIV-positive patients in Indonesia. *J Int AIDS Soc* 2014; **17**: 18821.
58. Roberts T. Initial Perspectives on the Implementation of the Cryptococcal Antigen Lateral Flow Assay at the Point-of-Care. African Society for Lab Medicine; 2014; Cape Town, South Africa; 2014.

59. Kapoor SW, Magambo KA, Kalluvya SE, Fitzgerald DW, Peck RN, Downs JA. Six-month outcomes of HIV-infected patients given short-course fluconazole therapy for asymptomatic cryptococcal antigenemia. *AIDS* 2015; **29**(18): 2473-8.
60. Govender NP, Roy M, Mendes JF, Zulu TG, Chiller TM, Karstaedt AS. Evaluation of screening and treatment of cryptococcal antigenaemia among HIV-infected persons in Soweto, South Africa. *HIV Med* 2015; **16**(8): 468-76.
61. Mfinanga S, Chanda D, Kivuyo SL, et al. Cryptococcal meningitis screening and community-based early adherence support in people with advanced HIV infection starting antiretroviral therapy in Tanzania and Zambia: an open-label, randomised controlled trial. *Lancet* 2015; **385**(9983): 2173-82.
62. Pac L, Horwitz MM, Namutebi AM, et al. Implementation and operational research: Integrated pre-antiretroviral therapy screening and treatment for tuberculosis and cryptococcal antigenemia. *J Acquir Immune Defic Syndr* 2015; **68**(5): e69-76.
63. Morawski BM, Boulware DR, Nalintya E, et al. Pre-ART cryptococcal antigen titer associated with preemptive fluconazole failure. Abstract 159. Conference on Retroviruses and Opportunistic Infections. Boston; 2016.
64. Longley N, Jarvis JN, Meintjes G, et al. Cryptococcal antigen screening in patients initiating ART in South Africa: A prospective cohort study. *Clin Infect Dis* 2016; **62**(5): 581-7.
65. Oladele RO, Akanmu AS, Nwosu AO, Ogunsola FT, Richardson MD, Denning DW. Cryptococcal Antigenemia in Nigerian Patients With Advanced Human Immunodeficiency Virus: Influence of Antiretroviral Therapy Adherence. *Open Forum Infect Dis* 2016; **3**(2): ofw055.
66. Ezeanolue EE, Nwizu C, Greene GS, et al. Brief Report: Geographical Variation in Prevalence of Cryptococcal Antigenemia Among HIV-Infected, Treatment-Naïve Patients in Nigeria: A Multicenter Cross-Sectional Study. *J Acquir Immune Defic Syndr* 2016; **73**(1): 117-21.
67. Vallabhaneni S, Longley N, Smith M, et al. Implementation and Operational Research: Evaluation of a Public-Sector, Provider-Initiated Cryptococcal Antigen Screening and Treatment Program, Western Cape, South Africa. *J Acquir Immune Defic Syndr* 2016; **72**(2): e37-42.
68. Ogouyemi-Hounto A, Zannou DM, Ayihounton G, et al. [Prevalence and factors associated with cryptococcal antigenemia in HIV-infected patients in Cotonou/Benin]. *J Mycol Med* 2016; **26**(4): 391-7.
69. Kadam D, Chandanwale A, Bharadwaj R, et al. High prevalence of cryptococcal antigenaemia amongst asymptomatic advanced HIV patients in Pune, India. *Indian J Med Microbiol* 2017; **35**(1): 105-8.
70. Grant AD, Djomand G, De Cock KM. Natural history and spectrum of disease in adults with HIV/AIDS in Africa. *AIDS* 1997; **11 Suppl B**: S43-54.
71. Oumar AA, Dao S, Ba M, Poudiougou B, Diallo A. [Epidemiological, clinical and prognostic aspects of cryptococcal meningitis in hospital area of Bamako, Mali]. *Rev Med Brux* 2008; **29**(3): 149-52.
72. Wajanga BM, Kalluvya S, Downs JA, Johnson WD, Fitzgerald DW, Peck RN. Universal screening of Tanzanian HIV-infected adult inpatients with the serum cryptococcal antigen to improve diagnosis and reduce mortality: an operational study. *J Int AIDS Soc* 2011; **14**: 48.

73. Apetse K, Assogba K, Kevi K, Balogou AA, Pitche P, Grunitzky E. [Opportunistic infections of the HIV/AIDS in adults in hospital settings in Togo]. *Bull Soc Pathol Exot* 2011; **104**(5): 352-4.
74. Oyella J, Meya D, Bajunirwe F, Kanya MR. Prevalence and factors associated with cryptococcal antigenemia among severely immunosuppressed HIV-infected adults in Uganda: a cross-sectional study. *J Int AIDS Soc* 2012; **15**(1): 15.
75. Dzoyem JP, Kechia FA, Ngaba GP, Lunga PK, Lohoue PJ. Prevalence of cryptococcosis among HIV-infected patients in Yaounde, Cameroon. *Afr Health Sci* 2012; **12**(2): 129-33.
76. Luma HN, Temfack E, Halle MP, Tchaleu BC, Mapoure YN, Koulla-Shiro S. Cryptococcal meningoencephalitis in human immunodeficiency virus/acquired immunodeficiency syndrome in douala, cameroon: a cross sectional study. *N Am J Med Sci* 2013; **5**(8): 486-91.
77. Okome-Nkoumou M, Guiyedi V, Ondounda M, et al. Opportunistic Diseases in HIV-Infected Patients in Gabon following the Administration of Highly Active Antiretroviral Therapy: A Retrospective Study. *The American Journal of Tropical Medicine and Hygiene* 2014; **90**(2): 211-5.
78. Katchanov J, Jefferys L, Tominski D, et al. Cryptococcosis in HIV-infected hospitalized patients in Germany: Evidence for routine antigen testing. *J Infect* 2015; **71**(1): 110-6.
79. Frola C, Guelfand L, Socias E, Sued O, Perez H, Canh P. Prevalencia del antígeno de *Cryptococcus* y valoración de su detección como método de tamizaje en pacientes con infección avanzada por VIH. XVII Congreso de la Asociación Panamericana de Infectología; 2015 16–19 May 2015; Quito, Ecuador; 2015.
80. Vidal JE, Toniolo C, Paulino A, et al. Asymptomatic cryptococcal antigen prevalence detected by lateral flow assay in hospitalised HIV-infected patients in Sao Paulo, Brazil. *Trop Med Int Health* 2016; **21**(12): 1539-44.